

Clinical Policy: Belinostat (Beleodaq)

Reference Number: CP.PHAR.311 Effective Date: 02.01.17 Last Review Date: 11.22 Line of Business: HIM, Medicaid

Coding Implications Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

Description

Belinostat (Beleodaq[®]) is a histone deacetylase inhibitor.

FDA Approved Indication(s)

Beleodaq is indicated for the treatment of patients with relapsed or refractory peripheral T-cell lymphoma (PTCL).

This indication is approved under accelerated approval based on tumor response rate and duration of response. An improvement in survival or disease-related symptoms has not been established. Continued approval for this indication may be contingent upon verification and description of clinical benefit in the confirmatory trial.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Beleodaq is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Peripheral T-Cell Lymphoma (must meet all):

- 1. Diagnosis of PTCL (see Appendix D for examples of PTCL subtypes);
- 2. Prescribed by or in consultation with an oncologist or hematologist;
- 3. Age \geq 18 years;
- 4. Request meets one of the following (a or b):*
 - a. Dose does not exceed $1,000 \text{ mg/m}^2$ per day on days 1-5 of a 21-day cycle;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration: 6 months

B. NCCN-Recommended Off-Label Indications (must meet all):

- 1. Diagnosis of one of the following (a, b, c, d, or e):
 - a. Primary cutaneous anaplastic large cell lymphoma (ALCL) with multifocal lesions, or cutaneous ALCL with regional nodes;
 - b. Adult T-cell leukemia/lymphoma;
 - c. Extranodal NK/T-cell lymphoma, nasal type;

CLINICAL POLICY Belinostat



- d. Hepatosplenic gamma-delta T-cell lymphoma;
- e. Breast implant ALCL;
- 2. Prescribed by or in consultation with an oncologist or hematologist;
- 3. Age \geq 18 years;

*Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration: 6 months

C. Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: HIM.PA.33 for health insurance marketplace and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: HIM.PA.103 for health insurance marketplace and CP.PMN.16 for Medicaid; or
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: HIM.PA.154 for health insurance marketplace and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. All Indications in Section I (must meet all):

- 1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Beleodaq for a covered indication and has received this medication for at least 30 days;
- 2. Member is responding positively to therapy;
- 3. If request is for a dose increase, request meets one of the following (a or b):*
 - a. New dose does not exceed 1,000 mg/m² per day on days 1-5 of a 21-day cycle;
 - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: HIM.PA.33 for health insurance marketplace and CP.PMN.255 for Medicaid; or

CLINICAL POLICY Belinostat



- b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: HIM.PA.103 for health insurance marketplace and CP.PMN.16 for Medicaid; or
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: HIM.PA.154 for health insurance marketplace and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – HIM.PA.154 for health insurance marketplace and CP.PMN.53 for Medicaid, or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key ALCL: anaplastic large cell lymphoma FDA: Food and Drug Administration

NCCN: National Comprehensive Cancer Network PTCL: peripheral T-cell lymphoma

Appendix B: Therapeutic Alternatives Not applicable

Appendix C: Contraindications/Boxed Warnings None reported

Appendix D: General Information

- PTCL subtypes/histologies:
 - PTCL, not otherwise specified;
 - Anaplastic large cell lymphoma;
 - Angioimmunoblastic T-cell lymphoma;
 - Enteropathy-associated T-cell lymphoma;
 - Monomorphic epitheliotropic intestinal T-cell lymphoma;
 - Nodal peripheral T-cell lymphoma with TFH phenotype;
 - Follicular T-cell lymphoma;

*PTLC is classified as a non-Hodgkin T-cell lymphoma. PTCL classification schemes are periodically advanced as new information becomes available; therefore, the above list is provided as general guidance. For additional information, see WHO's 2016 updated classification of hematological malignancies for a complete list of lymphoid neoplasms, including PTCL.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
PTCL	$1,000 \text{ mg/m}^2$ IV on days 1-5 of a 21-day cycle. Cycles can	1,000 mg/m ² /day
	be repeated every 21 days until disease progression or	
	unacceptable toxicity.	



VI. Product Availability

Single-dose vial: 500 mg

VII. References

- 1. Beleodaq Prescribing Information. Irvine, CA: Spectrum Pharmaceuticals, Inc.; April 2022. Available at: http://www.beleodaq.com/downloads/Beleodaq_PI.pdf. Accessed July 5, 2022.
- 2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug_compendium. Accessed July 5, 2022.
- 3. National Comprehensive Cancer Network. T-Cell Lymphomas Version 2.2022. Available at: https://www.nccn.org/professionals/physician_gls/pdf/t-cell.pdf. Accessed July 5, 2022.
- 4. National Comprehensive Cancer Network. Primary Cutaneous Lymphomas Version 2.2022. Available at: https://www.nccn.org/professionals/physician_gls/pdf/primary_cutaneous.pdf. Accessed July 5, 2022.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-todate sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J9032	Injection, belinostat, 10 mg

Reviews, Revisions, and Approvals	Date	P&T Approval Date
4Q 2018 annual review: added HIM-Medical Benefit line of business; summarized NCCN and FDA-approved uses for improved clarity; added specialist involvement in care; references reviewed and updated.	07.31.18	11.18
4Q 2019 annual review: added NCCN-recommended (with Category 2A or above) off-label uses: extranodal NK/T-cell lymphoma, nasal type, hepatosplenic gamma-delta T-cell lymphoma; references reviewed and updated.	08.14.19	11.19
4Q 2020 annual review: modified HIM-Medical Benefit to HIM line of business; added additional off-label indication cutaneous CD30+ T- cell lymphoma as per NCCN 2A or above off label indication; added Appendix D: PTCL subtypes per NCCN; references reviewed and updated.	08.14.20	11.20
4Q 2021 annual review: HIM.PHAR.21 changed to HIM.PA.154; references reviewed and updated.	06.30.21	11.21
4Q 2022 annual review: updated NCCN-recommended off-label uses: removed mycosis fungoides, cutaneous CD30+ T-cell lymphoma, and Sézary syndrome; added breast implant ALCL (Category 2A	07.05.22	11.22



Reviews, Revisions, and Approvals	Date	P&T Approval Date
recommendation); references reviewed and updated. Template changes applied to other diagnoses/indications.		

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

CLINICAL POLICY Belinostat



This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note: For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2017 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation.