

## **Clinical Policy: Crizotinib (Xalkori)**

Reference Number: CP.PHAR.90

Effective Date: 11.01.11

Last Review Date: 05.22

Line of Business: Commercial, HIM, Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### **Description**

Crizotinib (Xalkori<sup>®</sup>) is a kinase inhibitor.

### **FDA Approved Indication(s)**

Xalkori is indicated for the treatment of:

- Patients with metastatic non-small cell lung cancer (NSCLC) whose tumors are anaplastic lymphoma kinase (ALK) or ROS1-positive as detected by an FDA-approved test.
- Pediatric patients 1 year of age and older and young adults with relapsed or refractory, systemic anaplastic large cell lymphoma (ALCL) that is ALK-positive.
- Adult and pediatric patients 1 year of age and older with unresectable, recurrent, or refractory inflammatory myofibroblastic tumor (IMT) that is ALK-positive.

Limitation(s) of use: The safety and efficacy of Xalkori have not been established in older adults with relapsed or refractory, systemic ALK-positive ALCL.

### **Policy/Criteria**

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Xalkori is **medically necessary** when the following criteria are met:

#### **I. Initial Approval Criteria**

##### **A. Non-Small Cell Lung Cancer** (must meet all):

1. Diagnosis of recurrent, advanced or metastatic NSCLC;
2. Prescribed by or in consultation with an oncologist;
3. Age  $\geq$  18 years;
4. Disease is ALK, ROS1, MET exon 14 skipping, or high-level MET amplification positive;
5. Prescribed as a single agent;
6. For Xalkori requests, member must use generic crizotinib, if available, unless contraindicated or clinically significant adverse effects are experienced;
7. Request meets one of the following (a or b):\*
  - a. Dose does not exceed both of the following (i and ii):
    - i. 500 mg per day;
    - ii. 2 capsules per day;

- b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration:**

**Medicaid/HIM** – 6 months

**Commercial** – 12 months or duration of request, whichever is less

**B. Anaplastic Large Cell Lymphoma (must meet all):**

1. Diagnosis of relapsed or refractory ALCL (a peripheral T-cell lymphoma);
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age  $\geq$  1 year;
4. Disease is ALK-positive;
5. Prescribed as a single agent;
6. For Xalkori requests, member must use generic crizotinib, if available, unless contraindicated or clinically significant adverse effects are experienced;
7. Request meets one of the following (a or b):\*
  - a. Dose does not exceed both of the following (i and ii):
    - i. 560 mg/m<sup>2</sup> per day, up to a maximum of 1,000 mg per day;
    - ii. 4 capsules per day;
  - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration:**

**Medicaid/HIM** – 6 months

**Commercial** – 12 months or duration of request, whichever is less

**C. Inflammatory Myofibroblastic Tumor (must meet all):**

1. Diagnosis of IMT;
2. Prescribed by or in consultation with an oncologist;
3. Age  $\geq$  1 year;
4. Disease is ALK-positive;
5. Prescribed as a single agent;
6. For Xalkori requests, member must use generic crizotinib, if available, unless contraindicated or clinically significant adverse effects are experienced;
7. Request meets one of the following (a, b, or c):\*
  - a. For pediatric members, dose does not exceed both of the following (i and ii):
    - i. 560 mg/m<sup>2</sup> per day, up to a maximum of 1,000 mg per day;
    - ii. 4 capsules per day;
  - b. For adult members, dose does not exceed both of the following (i and ii):
    - i. 500 mg per day;
    - ii. 2 capsules per day;
  - c. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration:**

**Medicaid/HIM** – 6 months

**Commercial** – 12 months or duration of request, whichever is less

**D. Off-Label NCCN Compendium Recommended Indications** (must meet all):

1. Diagnosis of histiocytic neoplasm (Erdheim-Chester Disease, Langerhans Cell Histiocytosis, Rosai-Dorfman Disease);
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age  $\geq$  18 years;
4. Disease is ALK-positive;
5. Prescribed as a single agent;
6. For Xalkori requests, member must use generic crizotinib, if available, unless contraindicated or clinically significant adverse effects are experienced;
7. Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).\*

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration:**

**Medicaid/HIM** – 6 months

**Commercial** – 12 months or duration of request, whichever is less

**E. Other diagnoses/indications** (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

**II. Continued Therapy**

**A. All Indications in Section I** (must meet all):

1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Xalkori for a covered indication and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. For Xalkori requests, member must use generic crizotinib, if available, unless contraindicated or clinically significant adverse effects are experienced;

4. If request is for a dose increase, request meets one of the following (a, b, or c):\*
  - a. NSCLC and adult IMT: New dose does not exceed both of the following (i and ii):
    - i. 500 mg per day;
    - ii. 2 capsules per day;
  - b. ALCL and pediatric IMT: New dose does not exceed both of the following (i and ii):
    - i. 560 mg/m<sup>2</sup> per day, up to a maximum of 1,000 mg per day;
    - ii. 4 capsules per day;
  - c. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).

\*Prescribed regimen must be FDA-approved or recommended by NCCN

**Approval duration:**

**Medicaid/HIM** – 12 months

**Commercial** – 12 months or duration of request, whichever is less

**B. Other diagnoses/indications (must meet 1 or 2):**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid or evidence of coverage documents.

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

ALK: anaplastic lymphoma kinase

ALCL: anaplastic large cell lymphoma

FDA: Food and Drug Administration

IMT: inflammatory myofibroblastic tumor

MET: mesenchymal-epithelial transition

NCCN: National Comprehensive Cancer  
Network

NSCLC: non-small cell lung cancer

ROS1: ROS proto-oncogene 1

*Appendix B: Therapeutic Alternatives*

Not applicable

*Appendix C: Contraindications/Boxed Warnings*

None reported

*Appendix D: General Information*

- MET amplification is an oncogenic driver occurring in 1% to 5% of NSCLCs that confers a poor prognosis and lacks approved targeted therapies. High-level MET amplification is an emerging biomarker to identify novel therapies for patients with metastatic NSCL per NCCN. The definition of high-level MET amplification is evolving and may differ according to the assay used for testing. For NGS-based results, a copy number greater than 10 is consistent with high-level MET amplification. (NCCN NSCLC Guideline Version 1.2022)

**V. Dosage and Administration**

Indication	Dosing Regimen	Maximum Dose
NSCLC	250 mg PO BID	500 mg/day
ALCL	280 mg/m <sup>2</sup> PO BID	1,000 mg/day
IMT	Pediatric: 280 mg/m <sup>2</sup> PO BID Adult: 250 mg PO BID	Pediatric: 1,000 mg/day Adult: 500 mg/day

**VI. Product Availability**

Capsules: 200 mg, 250 mg

**VII. References**

1. Xalkori Prescribing Information. New York, NY: Pfizer, Inc.; July 2022. Available at <http://labeling.pfizer.com/showlabeling.aspx?id=676>. Accessed July 25, 2022.
2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at [www.nccn.org](http://www.nccn.org). Accessed February 14, 2022.
3. National Comprehensive Cancer Network Guidelines. Non-Small Cell Lung Cancer Version 1.2022. Available at [www.nccn.org](http://www.nccn.org). Accessed February 14, 2022.
4. National Comprehensive Cancer Network Guidelines. Histiocytic Neoplasms Version 2.2021. Available at [www.nccn.org](http://www.nccn.org). Accessed February 14, 2022.
5. National Comprehensive Cancer Network Guidelines. Soft Tissue Sarcoma Version 2.2022. Available at [www.nccn.org](http://www.nccn.org). Accessed July 25, 2022.
6. National Comprehensive Cancer Network Guidelines. T-Cell Lymphomas Version 1.2022. Available at [www.nccn.org](http://www.nccn.org). Accessed February 14, 2022.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
2Q 2018 annual review: policies combined for Commercial and Medicaid; added HIM line of business; age added; minimum dose	02.13.18	05.18

Reviews, Revisions, and Approvals	Date	P&T Approval Date
removed; off-label NSCLC recurrent disease added; off-label ALCL added; summarized NCCN and FDA approved uses for improved clarity; added specialist involvement in care; added continuity of care statement; references reviewed and updated.		
2Q 2019 annual review: NCCN designation of advanced added to NSCLC; references reviewed and updated.	02.19.19	05.19
2Q 2020 annual review: revised continued approval duration from 6 to 12 months; references reviewed and updated.	02.11.20	05.20
2Q 2021 annual review: RT4: updated with FDA-approved indication for ALCL (previously included as an NCCN supported off-label use) with age 1 year or older and dosing limits per label; oral oncology generic redirection language added; revised reference to HIM off-label use policy from HIM.PHAR.21 to HIM.PA.154; references reviewed and updated.	01.12.21	05.21
2Q 2022 annual review: for NSCLC, clarified criteria as MET as exon 14 skipping or high-level MET amplification positive per NCCN; added hematologist as specialist in ALCL; added criterion for Xalkori single-agent therapy for NSCLC, ALCL, and inflammatory myofibroblastic tumor per NCCN; added histiocytic neoplasms indications per NCCN category 2A; WCG.CP.PHAR.90 was retired and initial approval duration was consolidated to 6 months; Commercial approval durations revised from “Length of Benefit” to “12 months or duration of request, whichever is less”; references reviewed and updated.	02.14.22	05.22
RT4: updated criteria for new FDA approved indication of IMT and removed previous designation as off-label.	07.27.22	
Template changes applied to other diagnoses/indications.	10.12.22	

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering

benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note:**

**For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2011 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene<sup>®</sup> and Centene Corporation<sup>®</sup> are registered trademarks exclusively owned by Centene Corporation.