

Clinical Policy: Glucagon-Like Peptide-1 (GLP-1) Receptor Agonists

Reference Number: DE.PMN.183

Effective Date: 01.23

Last Review Date: 01.23

Line of Business: Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

The following agents contain a synthetic glucagon-like peptide-1 (GLP-1) receptor agonist and require prior authorization: dulaglutide (Trulicity[®]), exenatide ER (Bydureon[®], Bydureon BCise[®]), exenatide IR (Byetta[®]), liraglutide (Victoza[®]), liraglutide/insulin degludec (Xultophy[®]), lixisenatide (Adlyxin[®]), semaglutide (Ozempic[®], Rybelsus[®]), and tirzepatide* (Mounjaro[™]).

*Tirzepatide is a combination GLP-1 and glucose-dependent insulinotropic polypeptide (GIP) receptor agonist.

Requests for Soliqua[®] (lixisenatide/insulin glargine) should be reviewed against CP.PST.01 – Step Therapy Criteria.

FDA Approved Indication(s)

GLP-1 receptor agonists are indicated as adjunct to diet and exercise to improve glycemic control with type 2 diabetes mellitus. Bydureon, Bydureon BCise, and Victoza are indicated in patients 10 years of age and older, while the other GLP-1 receptor agonists are indicated in adults.

Ozempic, Trulicity, and Victoza are also indicated to reduce the risk of major adverse cardiovascular events in adults with type 2 diabetes mellitus and:

- Established cardiovascular disease (*Ozempic, Trulicity, Victoza*);
- Cardiovascular risk factors (*Trulicity only*).

Limitation(s) of use:

- Bydureon, Bydureon BCise, Xultophy, and Rybelsus are not recommended as a first-line therapy for patients inadequately controlled on diet and exercise.
- GLP-1 receptor agonists should not be used for the treatment of type 1 diabetes. Xultophy is not for the treatment of diabetic ketoacidosis.
- Xultophy has not been studied in combination with prandial insulin. In addition, they are not recommended for use in combination with any other product containing a GLP-1 receptor agonist.
- Other than Victoza and Xultophy, GLP-1 receptor agonists have not been studied in patients with a history of pancreatitis. Other antidiabetic therapies should be considered.
- Trulicity is not for patients with pre-existing severe gastrointestinal disease.
- Adlyxin is not recommended in patients with gastroparesis.
- Bydureon and Bydureon BCise are extended-release formulations of exenatide. Do not coadminister with other exenatide containing products.
- Victoza and Xultophy contain liraglutide and should not be co-administered with other liraglutide-containing products.

CLINICAL POLICY

Glucagon-Like Peptide 1 (GLP-1) Receptor Agonists

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation® that GLP-1 receptor agonists are **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Type 2 Diabetes Mellitus (must meet all):

1. Diagnosis of type 2 diabetes mellitus;
2. Age is one of the following (a or b):
 - a. Bydureon, Bydureon BCise, Victoza: ≥ 10 years;
 - b. All other GLP-1 receptor agonists: ≥ 18 years;
3. Member meets one of the following (a or b):
 - a. Failure of ≥ 3 consecutive months of metformin as evidenced by HbA1c $\geq 7\%$, unless contraindicated or clinically significant adverse effects are experienced;
 - b. For antidiabetic medication-naïve members, requested agent is approvable if intended for concurrent use with metformin due to HbA1c $\geq 8.5\%$ (drawn within the past 3 months);
4. Failure of ≥ 3 consecutive month trial of a sodium-glucose co-transporter 2 (SGLT2) inhibitor or SGLT2 inhibitor-containing product (see *Appendix B*), unless clinically significant adverse effects are experienced or all are contraindicated;
5. If request is for a non-preferred GLP-1 receptor agonist, failure of ≥ 3 consecutive months of a preferred GLP-1 receptor agonist, unless (a or b):
 - a. Clinically significant adverse effects are experienced or all are contraindicated;
 - b. Request is for Ozempic, Trulicity, or Victoza, and member has established cardiovascular disease (e.g., ASCVD) or multiple cardiovascular risk factors;
6. Dose does not exceed the FDA-approved maximum recommended dose (see *Section V*).

Approval duration: 12 months

B. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Type 2 Diabetes Mellitus (must meet all):

1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy;
3. If request is for a dose increase, new dose does not exceed the FDA-approved maximum recommended dose (see *Section V*).

Approval duration: 12 months

CLINICAL POLICY

Glucagon-Like Peptide 1 (GLP-1) Receptor Agonists

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.
Approval duration: Duration of request or 12 months (whichever is less); or
2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

AACE: American Association of Clinical Endocrinologists

ACE: American College of Endocrinology

ADA: American Diabetes Association

ASCVD: atherosclerotic cardiovascular disease

ER: extended-release

FDA: Food and Drug Administration

GIP: glucose-dependent insulinotropic polypeptide

GLP-1: glucagon-like peptide-1

HbA1c: glycated hemoglobin

IR: immediate-release

SGLT2: sodium-glucose co-transporter 2

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
metformin (Fortamet [®] , Glucophage [®] , Glucophage XR, Glumetza [®])	Regular-release (Glucophage): 500 mg PO BID or 850 mg PO QD; increase as needed in increments of 500 mg/week or 850 mg every 2 weeks Extended-release: <ul style="list-style-type: none"> • Fortamet, Glumetza: 1,000 mg PO QD; increase as needed in increments of 500 mg/week • Glucophage XR: 500 mg PO QD; increase as needed in increments of 500 mg/week 	Regular-release: 2,550 mg/day Extended-release: 2,000 mg/day
SGLT2 Inhibitors		
Farxiga [®] (dapagliflozin)	5 mg PO QD	10 mg/day

CLINICAL POLICY

Glucagon-Like Peptide 1 (GLP-1) Receptor Agonists

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
	To reduce the risk of hospitalization for heart failure, the recommended dose is 10 mg PO QD	
Glyxambi [®] (empagliflozin/linagliptin)	One 10/5 mg tablet PO QD	25/5 mg/day
Invokamet [®] (canagliflozin/metformin)	One 50/500 mg tablet PO BID	300/2,000 mg/day
Invokamet [®] XR (canagliflozin/metformin)	Two 50/500 mg tablets PO QD	300/2,000 mg/day
Invokana [®] (canagliflozin)	100 mg PO QD	300 mg/day
Jardiance [®] (empagliflozin)	10 mg PO QD	25 mg/day
Qtern [®] (dapagliflozin/saxagliptin)	One 5/5 mg tablet PO QD	10/5 mg/day
Segluromet [™] (ertugliflozin/ metformin)	Individualized dose PO BID	15/2,000 mg/day
Steglatro [™] (ertugliflozin)	5 mg PO QD	15 mg/day
Steglujan [™] (ertugliflozin/sitagliptin)	One 5/100 mg tablet PO QD	15/100 mg/day
Synjardy [®] (empagliflozin/metformin)	Individualized dose PO BID	25/2,000 mg/day
Synjardy [®] XR (empagliflozin/metformin)	Individualized dose PO QD	25/2,000 mg/day
Trijardy [™] XR (empagliflozin/linagliptin/ metformin)	Individualized dose PO QD	25/5/2,000 mg/day
Xigduo [®] XR (dapagliflozin/metformin)	Individualized dose PO QD	10/2,000 mg/day

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s):
 - Hypersensitivity to any product components
 - Personal or family history of medullary thyroid carcinoma or multiple endocrine neoplasia syndrome type 2 (*all GLP-1 receptor agonists other than Byetta, and Adlyxin*)
 - Use during episodes of hypoglycemia (*Xultophy only*)
 - History of drug-induced immune-mediated thrombocytopenia from exenatide products (*Bydureon, Bydureon BCise, and Byetta only*)
- Boxed warning(s): thyroid C-cell tumors (*all GLP-1 receptor agonists other than Byetta, and Adlyxin*)

CLINICAL POLICY

Glucagon-Like Peptide 1 (GLP-1) Receptor Agonists

Appendix D: General Information

- Per the American Diabetes Association (ADA) and American Association of Clinical Endocrinologists and American College of Endocrinology (AACE/ACE) guidelines:
 - Metformin is recommended for all patients with type 2 diabetes. Monotherapy is recommended for most patients; however:
 - Starting with dual therapy (i.e., metformin plus another agent, such as a sulfonylurea, thiazolidinedione, dipeptidyl peptidase-4 inhibitor, sodium-glucose co-transporter inhibitor, GLP-1 receptor agonist, or basal insulin) may be considered for patients with baseline HbA1c $\geq 1.5\%$ above their target per the ADA ($\geq 7.5\%$ per the AACE/ACE). According to the ADA, a reasonable HbA1c target for many non-pregnant adults is $< 7\%$ ($\leq 6.5\%$ per the AACE/ACE).
 - Starting with combination therapy with insulin may be considered for patients with baseline HbA1c $> 10\%$ per the ADA ($> 9\%$ if symptoms are present per the AACE/ACE).
 - If the target HbA1c is not achieved after approximately 3 months of monotherapy, dual therapy should be initiated. If dual therapy is inadequate after 3 months, triple therapy should be initiated. Finally, if triple therapy fails to bring a patient to goal, combination therapy with insulin should be initiated. Each non-insulin agent added to initial therapy can lower HbA1c by 0.7-1%.
- Although Trulicity is currently the only GLP-1 receptor agonist that is FDA approved for use in patients with multiple cardiovascular risk factors, the ADA guidelines recognize Ozempic, Trulicity, and Victoza as agents that confer cardiovascular benefit and recommend the use of any of the three in patients at high risk of ASCVD, without preference for any one over the other. In addition, patients with multiple cardiovascular risk factors were included in each drug's cardiovascular outcomes trial.
- Examples of cardiovascular risk factors may include but are not limited to: dyslipidemia, hypertension, obesity, a family history of premature coronary disease, smoking, chronic kidney disease, and presence of albuminuria.
- According to the ADA, ASCVD includes coronary heart disease, cerebrovascular disease, or peripheral arterial disease presumed to be of atherosclerotic origin. Indicators of high ASCVD risk are age ≥ 65 years with coronary, carotid, or lower-extremity artery stenosis $> 50\%$ or left ventricular hypertrophy.

V. Dosage and Administration

Drug Name	Dosing Regimen	Maximum Dose
Adlyxin (lixisenatide)	Initial dose: 10 mcg SC QD for 14 days Maintenance dose: 20 mcg SC QD	20 mcg/day
Bydureon (exenatide ER)	2 mg SC once weekly	2 mg/week
Bydureon BCise (exenatide ER)	2 mg SC once weekly	2 mg/week
Byetta (exenatide IR)	5 mcg to 10 mcg SC BID	20 mcg/day
Mounjaro (tirzepatide)	Initial dose: 2.5 mg SC once weekly. May increase by 2.5 mg every 4 weeks up to 15 mg once weekly	15 mg/week

CLINICAL POLICY

Glucagon-Like Peptide 1 (GLP-1) Receptor Agonists

Drug Name	Dosing Regimen	Maximum Dose
Ozempic (semaglutide)	0.25 mg to 2 mg SC once weekly, increased no more frequently than every 4 weeks	2 mg/week
Rybelsus (semaglutide)	Initial dose: 3 mg PO QD. After 30 days on the 3 mg dose, increase to 7 mg PO QD. May increase to 14 mg PO QD if needed after at least 30 days on the 7 mg dose	14 mg/day
Trulicity (dulaglutide)	0.75 mg to 1.5 mg SC once weekly. May increase to 3 mg once weekly if needed after at least 4 weeks on 1.5 mg dose. May further increase to 4.5 mg once weekly if needed after at least 4 weeks on 3 mg dose.	4.5 mg/week
Victoza (liraglutide)	Initial: 0.6 mg SC QD for 7 days Maintenance: 1.2 mg to 1.8 mg SC QD	1.8 mg/day
Xultophy (liraglutide/insulin degludec)	Treatment naïve to basal insulin or GLP-1 receptor agonist: 10 units (10 units of insulin/0.36 mg liraglutide) SC QD Treatment experienced to basal insulin or GLP-1 receptor agonist: 16 units (16 units insulin/0.58 mg liraglutide) SC QD	50 units insulin/1.8 mg liraglutide/day

VI. Product Availability

Drug Name	Availability
Adlyxin (lixisenatide)	Multi-dose prefilled pen: 50 mcg/mL in 3 mL (14 doses; 10 mcg/dose), 100 mcg/mL in 3 mL (14 doses; 20 mcg/dose)
Bydureon (exenatide ER)	<ul style="list-style-type: none"> Single-dose tray: 2 mg vial Single-dose prefilled pen: 2 mg pen
Bydureon BCise (exenatide ER)	Single-dose autoinjector: 2 mg
Byetta (exenatide IR)	Prefilled pen: 5 mcg/dose (0.02 mL) in 1.2 mL (60 doses), 10 mcg/dose (0.04 mL) in 2.4 mL (60 doses)
Mounjaro (tirzepatide)	Single-dose prefilled pen: 2.5 mg/0.5 mL, 5 mg/0.5 mL, 7.5 mg/0.5 mL, 10 mg/0.5 mL, 12.5 mg/0.5 mL, 15 mg/0.5 mL
Ozempic (semaglutide)	Prefilled pen: 2 mg/1.5 mL (1.34 mg/mL) for 0.25 mg dose (4 doses of 0.25 mg and 2 doses of 0.5 mg per pen) or 0.5 mg dose (4 doses per pen); 2 mg/1.5 mL (1.34 mg/mL) for 1 mg dose (2 doses per pen); 4 mg/3 mL (1.34 mg/mL) for 1 mg dose (4 doses per pen); 8 mg/3 mL (2.68 mg/mL) for 2 mg dose (4 doses per pen)
Rybelsus (semaglutide)	Tablets: 3 mg, 7 mg, 14 mg

CLINICAL POLICY

Glucagon-Like Peptide 1 (GLP-1) Receptor Agonists

Drug Name	Availability
Trulicity (dulaglutide)	Single-dose prefilled pen: 0.75 mg/0.5 mL, 1.5 mg/0.5 mL, 3 mg/0.5 mL, 4.5 mg/0.5 mL
Victoza (liraglutide)	Multi-dose prefilled pen: 18 mg/3 mL (6 mg/mL; delivers doses of 0.6 mg, 1.2 mg, or 1.8 mg)
Xultophy (liraglutide/insulin degludec)	Single-patient use pen: 3.6 mg/100 units per mL in 3 mL

VII. References

1. American Diabetes Association. Standards of medical care in diabetes—2022. *Diabetes Care*. 2022; 45(suppl 1): S1-S264. Updated December 2021. Accessed April 13, 2022.
2. Adlyxin Prescribing Information. Bridgewater, NJ: Sanofi-aventis US LLC; July 2021. Available at: www.adlyxin.com. Accessed September 16, 2021.
3. Bydureon Prescribing Information. Wilmington, DE: AstraZeneca Pharmaceuticals, LP; July 2021. Available at: www.bydureon.com. Accessed September 16, 2021.
4. Bydureon BCise Prescribing Information. Wilmington, DE: AstraZeneca Pharmaceuticals, LP; July 2021. Available at: www.bydureonbcise.com. Accessed September 16, 2021.
5. Byetta Prescribing Information. Wilmington, DE: AstraZeneca Pharmaceuticals, LP; June 2021. Available at: www.byetta.com. Accessed September 16, 2021.
6. Trulicity Prescribing Information. Indianapolis, IN: Eli Lilly and Company, Inc; April 2021. Available at: www.trulicity.com. Accessed September 16, 2021.
7. Victoza Prescribing Information. Princeton, NJ: Novo Nordisk Inc; November 2020. Available at: www.victoza.com. Accessed September 16, 2021.
8. Xultophy Prescribing Information. Bagsvaerd, Denmark: Novo Nordisk A/S; November 2019. Available at: www.xultophy.com. Accessed September 16, 2021.
9. Garber AJ, Handelsman Y, Grunberger G, et al. Consensus statement by the American Association of Clinical Endocrinologists and American College of Endocrinology on the comprehensive type 2 diabetes management algorithm – 2020 executive summary. *Endocr Pract*. 2020; 26(1): 107-139.
10. Ozempic Prescribing Information. Bagsvaerd, Denmark: Novo Nordisk A/S; March 2022. Available at: www.ozempic.com. Accessed April 13, 2022.
11. Rybelsus Prescribing Information. Bagsvaerd, Denmark: Novo Nordisk A/S; April 2021. Available at: www.rybelsuspro.com. Accessed September 16, 2021.
12. Mounjaro Prescribing Information. Indianapolis, IN: Lilly USA, LLC; May 2022. Available at: www.mounjaro.com. Accessed May 31, 2022.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy Created	01.23	01.23

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program

CLINICAL POLICY

Glucagon-Like Peptide 1 (GLP-1) Receptor Agonists

approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

CLINICAL POLICY

Glucagon-Like Peptide 1 (GLP-1) Receptor Agonists

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.