

Clinical Policy: Milnacipran (Savella)

Reference Number: DE.PMN.125

Effective Date: 01.23

Last Review Date: 01.23

Line of Business: Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Milnacipran (Savella[®]) is a selective serotonin and norepinephrine reuptake inhibitor (SNRI).

FDA Approved Indication(s)

Savella is indicated for the management of fibromyalgia. Savella is not approved for use in pediatric patients.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Savella is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Fibromyalgia (must meet all):

1. Diagnosis of fibromyalgia;
2. Age \geq 18 years;
3. Failure of an adequate trial of at least two preferred* FDA-approved drugs for the indication and/or drugs that are considered the standard of care within the same drug class on the PDL, when such agents exist, at maximum indicated doses, unless clinically significant adverse effect are experienced, or all are contraindicated;
 - a. * Generic is preferred, if available, and brand is not the preferred agent;
4. Dose does not exceed 200 mg (2 tablets) per day.

Approval duration:

Medicaid – 12 months

B. Depression (off-label) (must meet all):

1. Diagnosis of depression;
2. Age \geq 18 years;
3. Failure of an adequate trial of at least two preferred* FDA-approved drugs for the indication and/or drugs that are considered the standard of care within the same drug class on the PDL, when such agents exist, at maximum indicated doses, unless clinically significant adverse effect are experienced, or all are contraindicated;
 - * Generic is preferred, if available, and brand is not the preferred agent;
4. Dose does not exceed 200 mg (2 tablets) per day.

CLINICAL POLICY

Milnacipran

Approval duration:

Medicaid – 12 months

C. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53.

II. Continued Therapy

A. All Indications in Section I (must meet all):

1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy;
3. If request is for a dose increase, new dose does not exceed 200 mg (2 tablets) per day.

Approval duration:

Medicaid – 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.
Approval duration: Duration of request or 12 months (whichever is less); or
2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.PMN.53 or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

FDA: Food and Drug Administration

MAOI: monoamine oxidase inhibitor

SNRI: selective serotonin and norepinephrine reuptake inhibitor

SSRI: selective serotonin reuptake inhibitor

TCA: tricyclic antidepressant

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
amitriptyline (Elavil®)	Fibromyalgia: 10 mg to 50 mg PO once daily Depression: 50-100 mg PO daily	150 mg/day

CLINICAL POLICY

Milnacipran

nortriptyline (Pamelor [®])	Fibromyalgia: 25 mg to 50 mg PO once daily	150 mg/day
cyclobenzaprine (Flexeril [®])	Fibromyalgia: 10 mg PO every morning and 20 mg at bedtime	30 mg/day
bupropion (Wellbutrin [®])	Depression: 100 mg PO three times daily	450 mg/day
bupropion SR (Wellbutrin SR [®])	Depression: 150 mg PO twice daily	400 mg/day
bupropion XL (Wellbutrin XL [®])	Depression: 150 -300 mg PO once daily	450 mg/day
citalopram (Celexa [®])	Depression: 20-40 mg PO once daily	40 mg/day
desvenlafaxine succinate (Pristiq [®])	Depression: 50 mg PO once daily	50 mg/day
duloxetine (Cymbalta [®])	Fibromyalgia: 60 mg PO once daily Depression: 20 mg PO daily	60 mg/day
escitalopram (Lexapro [®])	Depression: 10 mg PO once daily	20 mg/day
fluoxetine (Prozac [®])	Depression: 20 mg PO once daily	80 mg/day
fluvoxamine (Luvox [®])	Depression (off-label): 50 mg PO once daily	300 mg/day
mirtazapine (Remeron [®])	Depression: 15 mg PO once daily	45 mg/day
paroxetine (Paxil [®])	Depression: 10 mg PO once daily	50 mg/day
paroxetine SR (Paxil CR [®])	Depression: 12.5 mg PO once daily	62.5 mg/day
sertraline (Zoloft [®])	Depression: 50 mg PO once daily	200 mg/day
venlafaxine(Effexor [®])	Depression: 75 mg PO once daily	375 mg/day
venlafaxine SR (Effexor XR [®])	Depression: 37.5 mg PO once daily	225 mg/day
doxepin (Sinequan [®])	Depression: 75 mg PO daily	300 mg/day
imipramine (Tofranil [®])	Depression: 75 mg PO daily	200 mg/day
trazodone (Desyrel [®])	Depression: 150mg PO in divided doses daily	400 mg/day

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): concomitant use or use within 14 days of discontinuing a monoamine oxidase inhibitor (MAOI) used to treat psychiatric disorders, use of an MAOI within 5 days of discontinuing Savella, initiation of Savella in patients currently treated with linezolid or IV methylene blue due to increased risk of serotonin syndrome.
- Boxed warning(s): increased risk of suicidal ideation, thinking, and behavior in children, adolescents, and young adults taking antidepressants for major depressive disorder (MDD) and other psychiatric disorders

Appendix D: General Information

- Class IIb recommendation in Micromedex for the treatment of depression.
- Use of MAOI with Savella concomitantly is contraindicated due to the risk of serious, sometimes, fatal, drug interactions with serotonergic drugs. These interactions have been associated with symptoms that include tremor, myoclonus, diaphoresis, nausea, vomiting,

CLINICAL POLICY

Milnacipran

flushing, dizziness, hyperthermia with features resembling neuroleptic malignant syndrome, seizures, rigidity, autonomic instability with possible rapid fluctuations of vital signs, and mental status changes that include extreme agitation progressing to delirium and coma. Allow at least 14 days after stopping an MAOI before starting Savella. Allow at least 5 days after stopping Savella before starting an MAOI.

- Savella should be stopped promptly, and linezolid or intravenous methylene blue can be administered. The patient should be monitored for symptoms of serotonin syndrome for 5 days or until 24 hours after the last dose of linezolid or intravenous methylene blue, whichever comes first. Therapy with Savella may be resumed 24 hours after the last dose of linezolid or intravenous methylene blue.
- Serotonin syndrome: Serotonin syndrome has been reported with SNRIs and SSRIs. Concomitant use of serotonergic drugs is not recommended.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Fibromyalgia	Based on efficacy and tolerability, PO dosing may be titrated according to the following schedule: <i>Day 1:</i> 12.5 mg once <i>Days 2-3:</i> 25 mg/day (12.5 mg twice daily) <i>Days 4-7:</i> 50 mg/day (25 mg twice daily) <i>After Day 7:</i> 100 mg/day (50 mg twice daily) Recommended dose is 100 mg/day PO (50 mg twice daily)	200 mg/day (100 mg twice daily)
Depression (off-label)	Initially, 12.5 to 25 mg PO twice daily. Based on individual response, the dose may be titrated to 100 mg PO twice daily	200 mg/day (100 mg twice daily)

VI. Product Availability

Tablets: 12.5 mg, 25 mg, 50 mg, 100 mg

VII. References

1. Savella Prescribing Information. Irvine, CA: Allergan USA, Inc.; September 2021. Available at: <https://www.savella.com/>. Accessed February 8, 2022.
2. Clauw DJ. Fibromyalgia: a clinical review. *JAMA*. 2014; 311(15): 1547-1555.
3. Häuser W, Walitt B, Fitzcharles M-A, Sommer C. Review of pharmacological therapies in fibromyalgia syndrome. *Arthritis Research & Therapy*. 2014;16(1):201. doi:10.1186/ar4441.
4. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc. Updated periodically. Accessed February 8, 2022.
5. American Psychiatric Association: Practice guideline for the treatment of patients with major depressive disorder 3rd edition. *Am J Psychiatry* 2010;167(suppl):1-152.
6. Kia S, Choy E. Update on treatment guideline in fibromyalgia syndrome with focus on pharmacology. *Biomedicines*. 2017,5,20;doi:10.3990
7. Micromedex® Healthcare Series [Internet database]. Greenwood Village, Colo: Thomson Healthcare. Updated periodically. Accessed February 8, 2022.

CLINICAL POLICY

Milnacipran

8. Heymann RE, Helfenstein M, Feldman D, et al. A double-blind, randomized, controlled study of amitriptyline, nortriptyline and placebo in patients with fibromyalgia. An analysis of outcome measures. Clin Exp Rheumatol. 2001;19(6)697-702.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created	11.22	01.23

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

CLINICAL POLICY

Milnacipran

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2012 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.