

Clinical Policy: Quetiapine Extended-Release (Seroquel XR)

Reference Number: CP.PMN.64

Effective Date: 09.01.15 Last Review Date: 02.22

Line of Business: Commercial, HIM, Medicaid Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

Description

Quetiapine extended-release (Seroquel XR®) is an atypical antipsychotic.

FDA Approved Indication(s)

Seroquel XR is indicated for the treatment of:

- Schizophrenia in adults and adolescents (13-17 years)
- Bipolar I disorder, manic or mixed episodes, in adults and children/adolescents (10-17 years)
- Bipolar disorder, depressive episodes, in adults
- Major depressive disorder (MDD), as adjunctive therapy with antidepressants, in adults

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation® that Seroquel XR is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Schizophrenia (must meet all):

- 1. Diagnosis of schizophrenia;
- 2. Age \geq 13 years;
- 3. Failure of a ≥ 4-week trial of quetiapine immediate-release (IR) at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;
- 4. Dose does not exceed 800 mg (2 tablets) per day.

Approval duration:

Medicaid/HIM – 12 months

Commercial – 12 months or duration of request, whichever is less

B. Bipolar Disorder (must meet all):

- 1. Diagnosis of bipolar disorder;
- 2. Age \geq 10 years;
- 3. Failure of a \geq 4-week trial of quetiapine IR at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;
- 4. Dose does not exceed 800 mg (2 tablets) per day.

Approval duration:

Medicaid/HIM – 12 months



Commercial – 12 months or duration of request, whichever is less

C. Major Depressive Disorder (must meet all):

- 1. Diagnosis of MDD;
- 2. Age \geq 18 years;
- 3. Failure of THREE antidepressants (e.g., selective serotonin reuptake inhibitor [SSRI], serotonin-norepinephrine reuptake inhibitor [SNRI], tricyclic antidepressant [TCA], bupropion, mirtazapine) from at least TWO different classes at up to maximally indicated doses, each used for ≥ 4 weeks, unless member is unable to satisfy this requirement due to clinically significant adverse effects experienced, member's age ≥ 65 years, or contraindication(s) to multiple antidepressants;
- 4. Failure of a ≥ 4-week trial of aripiprazole at up to maximally indicated doses, used concurrently with an antidepressant, unless contraindicated or clinically significant adverse effects are experienced;
- 5. Seroquel XR is prescribed concurrently with an antidepressant;
- 6. Dose does not exceed 300 mg (2 tablets) per day.

Approval duration:

Medicaid/HIM – 12 months

Commercial – 12 months or duration of request, whichever is less

D. Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. All Indications in Section I (must meet all):

- 1. Member meets one of the following (a, b, or c):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B);



- c. Documentation supports that member is currently receiving Seroquel XR for schizophrenia or bipolar disorder and has received this medication for at least 30 days;
- 2. Member is responding positively to therapy;
- 3. If request is for a dose increase, new dose does not exceed (a or b):
 - a. Schizophrenia, bipolar disorder: 800 mg (2 tablets) per day;
 - b. MDD: 300 mg (2 tablets) per day.

Approval duration:

Medicaid/HIM – 12 months

Commercial – 12 months or duration of request, whichever is less

B. Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business:
 CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid, or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

FDA: Food and Drug Administration

IR: immediate-release

MDD: major depressive disorder

SNRI: serotonin/norepinephrine

reuptake inhibitor

SSRI: selective serotonin reuptake inhibitor

TCA: tricyclic antidepressant

XR: extended-release



Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business

and may require prior authorization.

| and may require prior auth Drug Name | Dosing Regimen | Dose Limit/ Maximum Dose |
|---|--|--|
| Antipsychotics | | |
| quetiapine immediate-release (Seroquel®) | Schizophrenia Initial: 25 mg PO BID; target: 400 to 800 mg/day | 800 mg/day |
| | Bipolar Disorder Initial: 50 mg PO BID; target: 400 to 800 mg/day | |
| Selective Serotonin Reup | take Inhibitors (SSRIs) | |
| citalopram (Celexa®) | | 40 mg/day |
| escitalopram (Lexapro®) | | 20 mg/day |
| fluoxetine (Prozac®) | | Immediate-release: 80 mg/day (20 mg/day if pediatric) Delayed-release: 90 mg/week |
| fluvoxamine* | | 150 mg/day |
| (immediate-release) (Luvox®) | Major Depressive Disorder Refer to prescribing information | <i>y y</i> |
| paroxetine (Paxil®, Paxil CR®, Pexeva®) | | Immediate-release: 50 mg/day (40 mg/day if geriatric) Extended-release: 62.5 mg/day (50 mg/day if geriatric) |
| sertraline (Zoloft®) | | 200 mg/day (20 mg/day if age 6-11 years) |
| Serotonin-Norepinephrin | e Reuptake Inhibitors (SNRIs) | |
| desvenlafaxine (Pristiq®) duloxetine (Cymbalta®) | _ | 400 mg/day 120 mg/day |
| Fetzima [®] | Major Depressive Disorder | 120 mg/day |
| (levomilnacipran) | Refer to prescribing information | 120 mg day |
| venlafaxine (Effexor®, | | Extended-release: |
| Effexor XR®) | | 225 mg/day |
| Tricyclic Antidepressant | (TCAs) | 223 mg, day |
| amitriptyline (Elavil®) | | 150 mg/day |
| amoxapine | Major Depressive Disorder Refer to prescribing information | 400 mg/day (300 mg/day if geriatric) |



| Drug Name | Dosing Regimen Dose Limit/ | |
|--|----------------------------------|----------------------|
| | | Maximum Dose |
| clomipramine* | | 250 mg/day (200 |
| (Anafranil®) | | mg/day if pediatric) |
| desipramine | | 300 mg/day (100 |
| (Norpramin®) | | mg/day if pediatric) |
| doxepin (Sinequan®) | | 300 mg/day |
| imipramine HCl | | 200 mg/day (150 |
| (Tofranil [®]) | | mg/day if geriatric |
| | | or pediatric) |
| imipramine pamoate | | 200 mg/day (100 |
| (Tofranil PM®) | | mg/day if geriatric |
| | | or pediatric) |
| nortriptyline (Pamelor®) | | 150 mg/day |
| protriptyline (Vivactil®) | | 60 mg/day (30 |
| | | mg/day if geriatric |
| | | or pediatric) |
| trimipramine | | 200 mg/day (100 |
| (Surmontil®) | | mg/day if geriatric |
| | | or pediatric) |
| Monoamine Oxidase Inhi | bitors | T 11 |
| isocarboxazid | | 60 mg/day |
| (Marplan®) | | 00 /1 |
| phenelzine (Nardil®) | | 90 mg/day |
| selegiline (EMSAM® | Major Depressive Disorder | Transdermal: 12 |
| transdermal; Eldepryl®, | Refer to prescribing information | mg/24 hr |
| Zelapar [®] , Carbex [®]) | | Oral: 30 mg/day |
| tranylcypromine (Parnate®) | | 60 mg/day |
| Other Antidepressants | | |
| bupropion (Aplenzin®, | | Immediate-release: |
| Budeprion SR [®] , | | 450 mg/day (300 |
| Budeprion XL [®] , Forfivo | | mg/day if pediatric) |
| XL [®] , Wellbutrin [®] , | | Sustained-release: |
| Wellbutrin SR [®] , | | 400 mg/day |
| Wellbutrin XL®) | | Extended-release |
| (Venourin 7th) | | (HCl): 450 mg/day |
| | Major Depressive Disorder | Extended-release |
| | Refer to prescribing information | (HBr): 522 mg/day |
| mirtazapine (Remeron®) | | 45 mg/day |
| perphenazine/ | | 16 mg/day |
| amitriptyline (Triavil®) | | perphenazine and |
| | | 200 mg/day |
| | | amitriptyline |
| maprotiline (Ludiomil®) | | 150 mg/day |



| Drug Name | Dosing Regimen | Dose Limit/ |
|----------------------------|----------------|---------------------|
| | | Maximum Dose |
| nefazodone (Serzone®) | | 600 mg/day |
| trazodone (Desyrel®, | | Immediate-release: |
| Oleptro [®]) | | 400 mg/day |
| | | Extended-release: |
| | | 375 mg/day |
| vortioxetine (Trintellix®) | | 20 mg/day |
| vilazodone (Viibryd®) | | 40 mg/day |

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): known hypersensitivity to Seroquel XR or any components in the formulation
- Boxed warning(s): increased mortality in elderly patients with dementia-related psychosis; and suicidal thoughts and behaviors in children, adolescents, and young adults taking antidepressants.

V. Dosage and Administration

| Indication | Dosing Regimen | Maximum Dose |
|--------------------|---------------------------|---------------------------|
| Schizophrenia | Adults: | 800 mg/day |
| | Initial: 300 mg PO QD | |
| | Target: 400 to 800 mg/day | |
| | | |
| | Adolescents: | |
| | Initial: 50 mg PO QD | |
| | Target: 400 to 800 mg/day | |
| Bipolar I disorder | Manic or mixed episodes | Manic or mixed episodes |
| | Adults: | Adults: 800 mg/day |
| | Initial: 300 mg PO QD | Children and adolescents: |
| | Target: 400 to 800 mg/day | 600 mg/day |
| | Children and adolescents | |
| | Initial: 50 mg PO QD | Depressive episodes |
| | Target: 400 to 600 mg/day | 300 mg/day |
| | | |
| | Depressive episodes | |
| | Adults: | |
| | Initial: 50 mg PO QD | |
| | Target: 300 mg/day | |
| MDD | Adults: | 300 mg/day |
| | Initial: 50 mg PO QD | |
| | Target: 150 to 300 mg/day | |

VI. Product Availability

Extended-release tablets: 50 mg, 150 mg, 200 mg, 300 mg, 400 mg



VII. References

- 1. Seroquel XR Prescribing Information. Wilmington, DE: AstraZeneca Pharmaceuticals LP; September 2020. Available at: www.seroquelxr.com. Accessed November 13, 2021.
- 2. Lehman AF, Lieberman JA, Dixon LB, et al. Practice guideline for the treatment of patients with schizophrenia, second edition. Arlington, VA: American Psychiatric Association; February 2004. Available online at http://www.psych.org/psych pract/treatg/pg/prac guide.cfm. Accessed November 30, 2019.
- 3. Dixon L, Perkins D, Calmes C. Guideline watch: practice guideline for the treatment of patients with schizophrenia. Arlington, VA: American Psychiatric Association; September 2009. Available online at http://www.psych.org/psych_pract/treatg/pg/prac_guide.cfm. Accessed November 30, 2019.
- 4. Hirschfeld RMA, Bowden CL, Gitlin MJ, et al. Practice guideline for the treatment of patients with bipolar disorder, second edition. Arlington, VA: American Psychiatric Association; April 2002. Available online at http://www.psych.org/psych pract/treatg/pg/prac guide.cfm. Accessed November 30, 2019.
- 5. Hirschfeld RMA. Guideline watch: practice guideline for the treatment of patients with bipolar disorder. Arlington, VA: American Psychiatric Association; November 2005. Available online at http://www.psych.org/psych_pract/treatg/pg/prac_guide.cfm. Accessed November 30, 2019.
- 6. Gelenberg AJ, Freeman MP, Markowitz JC, et al. Practice guideline for the treatment of patients with major depressive disorder, third edition. Arlington, VA: American Psychiatric Association; May 2010. Available online at http://www.psych.org/psych_pract/treatg/pg/prac_guide.cfm. Accessed November 30, 2019.
- 7. Keepers G, Fochtmann L, Anzia J, et al. APA Practice guideline for the treatment of patients with schizophrenia, third edition. Am J Psychiatry. 2020 Sept;177(9):868-872.

| Reviews, Revisions, and Approvals | Date | P&T Approval |
|--|----------|-----------------|
| | | Date |
| 1Q18 annual review: Policies combined for Centene Medicaid, | 12.01.14 | 02.18 |
| Marketplace and Commercial lines of business; No significant | | |
| changes; References reviewed and updated | | |
| Medicaid: changed approval duration from 12 months to length of | 03.04.18 | 05.18 |
| benefit | | |
| 1Q 2019 annual review: no significant changes; references | 10.30.18 | 02.19 |
| reviewed and updated. | | |
| 1Q 2020 annual review: no significant changes; references | 11.30.19 | 02.20 |
| reviewed and updated. | | |
| Allowed members 65 years old or older to bypass redirections to | 03.27.20 | 08.20 |
| TCA for major depressive disorder. | | |
| 1Q 2021 annual review: no significant changes; references to | 11.29.20 | 02.21 |
| HIM.PHAR.21 revised to HIM.PA.154; references reviewed and | | |
| updated. | | |
| 1Q 2022 annual review: no significant changes; changed | 11.13.21 | 02.22 |
| Commercial line of business auth duration from Length of Benefit | | |



| Reviews, Revisions, and Approvals | Date | P&T Approval Date |
|--|----------|-------------------------|
| to 12 months or duration of request, whichever is less; references reviewed and updated. | | |
| Template changes applied to other diagnoses/indications and continued therapy section. | 10.10.22 | |

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.



Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2015 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.