

HEDIS® Pediatric Pocket Guide: 2026 Measurement Year

For a complete list of codes, please visit the NCQA website at [ncqa.org](https://www.ncqa.org), or see the HEDIS value sets. The following is a subset only of the NCQA approved codes.

All Well-Child Visits

Must include the following: Documentation of a visit with an acceptable provider type, the date of the visit, and services to validate a well-child visit was performed.

Measure	Codes	Best Practice
(W30) Well-Child Visits in the First 30 Months of Life (0–30 months) ^{1,2}	99381, 99382, 99391, 99392, 99461, S0302, Z00.110, Z00.111, Z00.121, Z00.129, Z00.2, Z02.84, Z76.1, Z76.2	<p>Well-Child Visits in the First 15 Months. For children who turned 15 months old during the measurement year: Six or more well-child visits.</p> <p>Well-Child Visits for 15–30 Months of Age. For children who turned 30 months of age during the measurement year: Two or more well-child visits.</p>
(WCV) Child and Adolescent Well-Care Visits (3–21 years of age) ^{1,2}	99382–99385, 99392–99395, S0302, S0610, S0612, S0613, Z00.00, Z00.01, Z00.121, Z00.129, Z00.2, Z00.3, Z01.411, Z01.419, Z02.5, Z02.84, Z76, Z76.2	One or more comprehensive well-care visits with a primary care provider (PCP) or obstetrician/gynecologist (OB/GYN) within the measurement year. Visits occurring anytime in the measurement year, including prior to or after the patient’s birthday, close the gap.

Respiratory Conditions

Measure	Codes	Best Practice
(CWP) Appropriate Testing for Pharyngitis ^{1,2}	<p>Group A Strep Test: 87070, 87071, 87081, 87430, 87650–87652, 87880</p>	<p>For episodes involving persons 3 years of age and older:</p> <ul style="list-style-type: none"> • Establish a <i>pharyngitis</i> diagnosis • Test for <i>group A streptococcus (strep)</i> • Treat with antibiotics <i>only if group A strep is detected</i> <p>Note: Perform a group A strep test before prescribing or dispensing antibiotics.</p>
	<p>Pharyngitis: J02.0, J02.8, J02.9, J03.00, J03.01, J03.80, J03.81, J03.90, J03.91</p>	

Weight Assessment and Counseling for Nutrition & Physical Activity

Measure	Codes	Best Practice
(WCC) Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (3–17 years of age) ^{1,2}	<p>BMI: Z68.51, Z68.52, Z68.53, Z68.54, Z68.55, Z68.56</p> <p>Nutrition Counseling: 97802–97804, G0270, G0271, G0447, S9449, S9452, S9470, Z71.3</p> <p>Physical Activity (PA): G0447, S9451, Z02.5, Z71.82</p>	Can be completed during a PCP or OB/GYN visit during the measurement year. Documentation must include height, weight, body mass index (BMI) percentile, nutrition behaviors and counseling, and current physical activity behaviors and counseling or guidance.

HEDIS® is a registered trademark of the National Committee for Quality Assurance (NCQA).

[delawarefirsthealth.com](https://www.delawarefirsthealth.com)

Lines of Business:

¹Medicaid

²Marketplace

Delaware First Health and Ambetter are affiliated products serving Medicaid and Health Insurance Marketplace members, respectively. The information presented here is representative of our network of products. If you have any questions, please contact Provider Relations.



Immunizations		
Measure	Codes	Best Practice
(CIS-E) Childhood Immunizations^{1,2} (Completed on or before the 2nd birthday)	90697, 90698, 90700, 90723	DTaP – 4 Doses
	90670, 90671, 90677, G0009	PCV – 4 Doses
	90697, 90698, 90713, 90723	IPV – 3 Doses
	90644, 90647, 90648, 90697, 90698, 90748	HiB – 3 Doses
	90697, 90723, 90740, 90744, 90747, 90748, G0010	Hep B – 3 Doses
	2 Doses: 90681 3 Doses: 90680	RV – 2 or 3 Doses
	90655–90658, 90660, 90661, 90672, 90674, 90685–90689, 90756	Flu – 2 Doses (LAIV meets criteria for one of the two required vaccinations if administered on the 2nd birthday)
	90710, 90716	VZV – 1 Dose
	90707, 90710	MMR – 1 Dose
90633	Hep A – 1 Dose	
(IMA-E) Immunizations for Adolescents^{1,2} (Completed on or before the 13th birthday)	90619, 90623, 90624, 90733, 90734	Meningococcal – 1 Dose
	90715	Tdap – 1 Dose
	90649–90651	HPV – 2 or 3 Doses

Lead Screening		
Measure	Codes	Best Practice
(LSC-E) Lead Screening in Children¹	83655	Percentage of children who had one or more capillary or venous lead blood tests by their 2nd birthday. At least one lead screening result documented by age 2.

(SDOH) Social Determinants of Health: Improving Patient Outcomes by Identifying and Addressing Social Needs		
Description	Codes	Best Practice
Occupational Exposure to Risk Factors	ICD-10: Z57.0–Z57.9	Include supplemental codes in the patient’s diagnosis section on a claim form
Problems Related to Education and Literacy	ICD-10: Z55.0–Z55.9	
Problems Related to Employment and Unemployment	ICD-10: Z56.0–Z56.9	
Problems Related to Physical Environment	ICD-10: Z58.0–Z58.9	
Problems Related to Housing And Economic Circumstances	ICD-10: Z59.0–Z59.9	
Problems Related to Social Environment	ICD-10: Z60.0–Z60.9	
Problems Related to Upbringing	ICD-10: Z62.0–Z62.9	
Problems Related to Primary Support Group, Including Family Circumstances	ICD-10: Z63.0–Z63.9	
Problems Related to Certain Psychosocial Circumstances	ICD-10: Z65.0–Z65.9	